

Neogen's 4-Step Rodent Control Program



Use Neogen's 4-step program to control rodents in and around your facilities.

Step 1. PREVENT:

Clean, clear, and implement preventative measures.

- a. Eliminate debris and other harborage for rodents in and around the facility.
- b. Remove easily accessible rodent food sources, including spilled feed.
 - Eliminating food sources increases the effectiveness of rodent bait.
- c. Clear/mow around the facility (~3 feet) to discourage rodents from entering.

Step 2. IDENTIFY:

Find the location and severity of the problem.

- a. Examinations done at dawn or dusk work best.
- b. Are there fresh droppings, burrows, tracks, or gnaw marks?
 - Be sure to inspect ceilings, other elevated runways, and facility grounds.
- c. Do you have rats, mice, or both?

Step 3. EXECUTE:

Implement baiting strategy according to target rodents and locations.

- a. Outside: Along the perimeter, place bait stations 15-30 feet apart for rats and 8-12 feet for mice. Use Havoc® XT Bloks, Havoc® Chunks, CyKill™ Blocks or Di-Kill™ Blocks and replace eaten bait at least weekly to start. Determine areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where rats or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.
- b. Inside: Use bait stations. Chunks or Blocks can be nailed or zip-stripped directly to overhead locations. Place packs should be placed in attics, ceilings, or other hard to access areas that rodents are using. Contact Neogen for more specific baiting suggestions.

Step 4. MONITOR:

Monitor the effectiveness of your program with record keeping. Frequency of baiting should change with increased activity.

This information is intended for use only as a guide in providing general information regarding the directions, warning and cautions associated with the use of this product. As with any pesticide, always follow the label instructions on the product package before using.



Neogen's Rotational Baiting Program

Better Bait. Better Results.



Neogen's Rotational Baiting strategy breaks the cycle of anticoagulant resistance.

CyKill™

2 Months
Bromethalin

Havoc®

4 Months
Brodifacoum

Di-Kill™

6 Months
Difenacoum



800/621-8829
859/254-1221

944 Nandino Blvd.
Lexington, KY 40511
inform@neogen.com
www.neogen.com