

Melons – see Cucurbits.

Onions (dry bulb & garlic) (1):

To control armyworm, cutworm, leafminer, onion maggot (adult), onion thrips, stinkbug. Do not exceed 6 applications per season.

Peppers (bell) (3):

To control beet armyworm, aphid, cabbage looper, corn earworm, European corn borer, flea beetle, pepper weevil, vegetable leafminer. Do not exceed 8 applications per season.

Potatoes (14):

To control beet armyworm, cabbage looper, Colorado potato beetle, cutworm, European corn borer, potato aphid, potato flea beetle, potato leafhopper, potato psyllid, potato tuberworm, tarnished plant bug. Do not exceed 8 applications per season.

Spinach (1):

To control aster leafhopper, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, leafhopper. Do not exceed 5 applications per season.

Sweet Corn (1):

To control corn earworm, corn rootworm (adult), European corn borer, fall armyworm, hop vine borer, leafhopper, southern armyworm, stalk borer. Dust every 5 days or as needed. Do not exceed 6 applications per season.

Tomatoes (0):

To control beet armyworm, cabbage looper, Colorado potato beetle, granulate cutworm, hornworm, vegetable leafminer, tomato fruitworm, tomato pinworm. Do not exceed 6 applications per season.

ORNAMENTALS

(including Roses, Flowers, Shrubs, and Small Trees)

Contact insects directly whenever possible.

Kills aphid, bagworms, cabbage looper, cicadas, corn earworms, cutworm, exposed thrips, fall cankerworm, fall webworm, inchworms, Japanese beetle (adults), leafhopper, leafminer, and omnivorous leafroller, mealybugs, spider mites, stinkbugs, adult strawberry root weevil, adult obscure root weevil, thrips and whitefly on the following plants:

African violets, Ageratum, Arizona Cypress, Aster, Azalea, Begonia, Camellia, Carnation, Chrysanthemum, Coleus, Common Ninebark, Common Snowberry, Crassula, Dahlias, Delphinium, Dogwood, Exacum, Euonymus, English Ivy, Fuchsia, Geranium, Gladiolus, Gold Bells, Grape Vines (ornamental), Hypoestes, Ivy, Laurel, Lilac, Mangold, Mock Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Poinsettia, Rhododendron, Rose, Snapdragon, Stocks, Taxus, Wax Plant, Wandering Jew, Weeping Fig, Zinnia.

FRUIT AND NUT TREES

The number in parentheses () following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest. Also includes relevant comments.

Apples (Do not apply after petal fall):

To control apple aphid, green fruitworm, redbanded leafroller, plum curculio, spotted tentiform leafminer, tarnished plant bug, white apple leafroller. Do not exceed 1 application per season.

Cherries (In the East, do not exceed 4 applications after petal fall):

To control green fruitworm, lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug. Do not exceed 6 applications per season.

Cherries (In the West, do not exceed 3 applications after petal fall):

To control green fruitworm, lesser peachtree borer, plum curculio, redbanded leafroller, rose chafer, tarnished plant bug. Do not exceed 4 applications per season.

Peaches (14):

To control green fruitworm, lesser peachtree borer, Oriental fruit moth, peach twig borer, plum curculio, rose chafer, tarnished plant bug. Dust lightly. Apply up to 5 times per season.

Pears – Dormant (May be applied dormant through the delayed dormant growth periods only): To control pear psylla. Do not exceed 2 applications per season.

Pears – Pre-bloom (May be applied from dormant through bud burst):

To control codling moth, green fruitworm, and pear psylla. Do not exceed 2 applications per season.

Walnuts (1):

To control codling moth, navel orangeworm, and walnut husk fly. Do not exceed 8 applications per season.

POULTRY

To control Northern Fowl Mites, Lice:

Apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area.

To control Northern Fowl Mites and Lice in Poultry Houses:

Apply to floors, roosts, and interior surfaces at a rate of 1 lb. per 40 sq. ft. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water.

PETS

USE ONLY ON DOGS OR CATS:

Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals.

To control Fleas, Ticks, Lice:

Use Prozap Garden & Poultry Dust only outside or in well-ventilated areas. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth, and genital areas. Rub or brush pet's hair to work dust onto the skin, paying close attention to legs and feet. Use ½ oz. on pets 20 pounds and under, and 1 oz. on pets over 20 pounds. Reapply every 2 weeks. Prozap Garden & Poultry Dust should not be applied by children. Do not use on puppies or kittens under 12 weeks of age. Dust pet bedding and doghouse using a shaker can or other dust applicator.

ANTS

To control Ants, Carpenter Ants:

Treat doors, around window frames, ant trails and hills, cracks and crevices, and other areas of entry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in a cool, dry, and secure area. Store in original sealed container away from food or feed.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling, if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DISCLAIMER

To the extent permitted by applicable law, Seller makes no representation or warranty, either express or implied, for results due to misuse, improper handling, or improper storage of this material. Nor does Seller assume any responsibility for injury to persons, crops, animals, soil, or property arising out of misuse, improper handling, or improper storage of this material.

For non-emergency (e.g., current product information), Call: 1-800-621-8289

Manufactured by:
Chem-Tech, Ltd.[™]
1006 Business Hwy 5
Pleasantville, IA 50225 U.S.A.

Prozap[®] is a Registered Trademark of Neogen[®] Corporation

Chem-Tech, Ltd.[™] is a wholly-owned subsidiary of Neogen[®] Corporation L6470-0419



NEOGEN

Item No. 1499540



NEOGEN

PROZAP[®] Garden & Poultry Dust

PERMETHRIN

⊕ Use on home garden fruits, vegetables and ornamental plants (including roses)

⊕ Controls Northern Fowl Mites on Poultry

⊕ Easy-to-use powder duster

Active Ingredient:

Permethrin.....0.25%

Other Ingredients.....99.75%

Total.....100.00%

* (3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloro-orethenyl)-2,2-dimethylcyclopropanecarboxylate
(Cis/Trans ratio: Min 35% (+/-) cis and max. 65% (+/-) trans.)

EPA Reg. No. 47000-149

EPA Est. No. 47000-IA-01

EPA Est. No. 89287-KS-2

EPA Est. No. 69208-MO-3

(First letter of batch number indicates establishment)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel and inside of leaflet for
First Aid and additional Precautionary Statements



Net Contents:
2 lbs.

FIRST AID	
If on Skin or on Clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for further treatment advice.
If in Eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.Remove contact lenses, if present after the first 15 minutes, then continue rinsing eye.Call a poison control center for further treatment advice.
Have product container or label with you when calling a poison control center, doctor, or going in for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas)	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

HUMANS: Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.
ANIMALS: Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash this pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Personal Protective Equipment (PPE):

Loaders, applicators, and handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes and socks;
- Chemical-resistant gloves made of any waterproof material, and
- NIOSH-approved respirator with: dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N, R, P, or HE filter.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Application with aerial or motorized ground equipment is prohibited.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions and precautions.

FOR PET USE: USE ONLY ON CATS AND DOGS.

USE RESTRICTIONS

- Do not breathe dust.
- Do not make applications in the rain.
- Do not water the treated area to the point of run-off. Use of hand-held power duster equipment is prohibited.
- Do not enter or allow others to enter until the dust has settled.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawn, turf, and listed vegetation; 3) Applications to building foundations, up to a maximum height of 3 feet.
- Do not water the treated area to the point of run-off.

HOME GARDENS

For use on home gardens. Not for use on plants being grown for commercial sale or commercial use.

Prozap Garden & Poultry Dust controls a wide range of insect pests. Apply when insects first appear before damage from feeding occurs. Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster, or other appropriate duster as available. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Allow dust to settle in treated areas before reentering. Do not allow dust to wind-drift to non-target plants or water supplies. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. For plants above chest height, consider using [brand insecticide].

VEGETABLES AND MELONS

The number in parentheses () following each crop grouping is the pre-harvest interval, e.g., the number of days that must be observed between application and harvest.

Artichokes (0): To control (kill) the artichoke plume moth and leafminer. Do not exceed 5 applications per season.

Asparagus (1):

To control the asparagus beetle and cutworm. Do not exceed 4 applications per season.

Asparagus (post-harvest):

To control the larval and adult stages of the asparagus beetle, tarnished plant bug, typhus bug, collards and adult stages of the Japanese beetle. Dust ferns or bush ground after spear harvest when insects are present. (Not for use in California.)

Celery – see leafy vegetables

Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage (tight-heading variation only), Chinese Broccoli (on Gal Lan, White Flowering) (1):

To control the beet armyworm, cabbage looper, cutworm, diamondback moth, flea beetle, imported cabbageworm, leafminer, plant bug, thrips. Apply every 5 days or as needed. Do not exceed 8 applications per season on cauliflower, Brussels sprouts and broccoli, or 5 applications on cabbage.

Corn – see sweet corn

Cucurbits (Balsam Pear, Bitter Melon, Chinese Wax gourd, Chinese Preserving Melon, Citron melon, Cucumber, Gherkin, Edible Gourds, Melons (including hybrids such as Cantaloupe, Casaba, Crenshaw, Honeydew, Honey Ball, Mango Melon, Muskmelon, Persian Melon), Pumpkin, Squash (Summer and Winter), Watermelon) (0):

To control aphid, cabbage looper, cucumber beetle (adult), cutworm, leafhopper, leafminer, melonworm, pickleworm, plant bug, squash bug. Do not exceed 8 applications per season.

Eggplant (3):

To control Colorado potato beetle, vegetable leafminer, flea beetle. Do not exceed 10 applications per season.

Leafy Vegetables (Amaranth, Arugula, Celery, Chervil, Corn Salad, Chrysanthemum (Edible-Leaved and Garland), Cress (Garden and Upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (Garden and Winter), Rhubarb, Spinach, and Swiss Chard) (1):

To control alfalfa looper, aphid, beet armyworm, cabbage looper, corn earworm, cutworm, European corn borer, fall armyworm, green cloverworm, southern armyworm, tobacco budworm, vegetable leafminer. Do not exceed 5 applications per season.