

RESTRICTED USE PESTICIDE

DUE TO HAZARD TO NON-TARGET SPECIES

For retail sale to and use only by Certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL

For Distribution and Use Only in the State of Oregon

Prozap[®] Zinc Phosphide Pellets

EPA Reg. No. 61282-49

EPA SLN No. OR-200008

**For Control of Voles and Mice in Clover Grown for Seed and
Only for Uses and Application Methods on this 24(c) SLN Label**

**This label valid until May 15, 2027, or until otherwise
amended, withdrawn, cancelled, or suspended.**

Active Ingredient: Zinc Phosphide.....	2.0%
<u>Other Ingredients</u>	<u>98.0%</u>
Total.....	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

(In addition to those on parent product (EPA Reg. No. 61282-49))

ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds, fish and other wildlife. Wildlife feeding on treated bait may be killed. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

To protect the Washington ground squirrel (*Urocyon washingtoni*), use is prohibited in shrub-steppe and grassland habitat with deep loam and sandy-loam soils of northern Gilliam, Morrow, and Umatilla Counties. In addition, use is prohibited on any lands that lie within 250 meters of occupied Washington ground squirrel sites. For information regarding these restrictions and to determine if the desired location of application lies within the scope of this restriction, please call the Oregon Department of Fish and Wildlife at 541-676-5230.

Users are advised that the Migratory Bird Treaty Act makes it unlawful to kill any migratory bird except as permitted by regulation. **Any bird deaths caused by application of this pesticide is a violation of both state and federal law.** In North America, all birds are considered migratory except for upland game birds (e.g. grouse, ptarmigans, prairie-chickens, quail, ring-necked pheasant and wild turkey) and introduced bird species (e.g. European starling, house sparrow, and rock dove). Refer to label booklet for Endangered Species Considerations.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This Section 24(c) SLN label and the Section 3 container and booklet labeling for Prozap® Zinc Phosphide Pellets (EPA Reg. No. 61282-49) must be in the possession of the user at the time of application. All applicable directions, restrictions, and precautions on the Section 3 container and booklet labeling and on this SLN label must be followed.

Use Restrictions: For control of voles (*Microtus spp.*), house mice, and deer mice in clover grown for seed as specified below.

Application Timing: Applications of this product under this SLN label may begin on **April 28, 2026**. Applications may be made through September 30, **2026**, or until Canada geese and cackling geese have been observed in the clover seed fields that were intended for treatment, whichever comes first. Depending on the year, Canada geese or cackling geese migrating southward may be observed in the field in mid-September or earlier. Above-ground applications of zinc phosphide may NOT be made if Canada geese or cackling geese have been observed in the field within the last fourteen (14) days, regardless of date.

Application Restrictions:

- Do not make applications if geese or other migratory birds have been observed in the field within the last fourteen (14) days.
- If geese are observed in the treated field within seven (7) days after broadcast application of this product, hazing is recommended. A hazing permit is not required.
- Keep out of any body of water. Do not apply where runoff is likely to occur.
- Do not graze animals on treated area or harvest any part of the treated crop for livestock feed.
- Do not apply if rain is forecast or irrigation is planned within the next three (3) days.
- Do not exceed four (4) applications per treated area per year.
- Anyone who applies bait by ground equipment must wear a dust mask and barrier laminate gloves. Applicators must wear barrier laminate gloves with hand baiting.

APPLICATION DIRECTIONS

For control of the vole complex (*Microtus spp.*), house mouse (*Mus musculus*), and deer mouse (*Peromyscus maniculatus*)

Broadcast Applications

When all restrictions indicated on this label are met, this product may be broadcast, either by air or by ground driven dispensing devices, cyclone seeder, or by hand in areas with heaviest vegetative cover. Apply at a rate of 6-10 pounds per acre. Use the higher label rate in areas with highest active population level. The number of vole holes is not an indicator of population level or activity.

Do not broadcast bait on bare ground. Do not apply by air when wind velocity exceeds 10 mph.

If applying this product after swathing the seed crop and while crop is drying in windrows, only apply between windrows. Do not apply over or on windrows or in such a manner that may result in zinc phosphide bait being combined with the clover. **This product may NOT be applied by air after swathing and while crop is drying in windrows.** However, aerial broadcast applications may resume after the seed crop has been combined and continue through **September 30, 2026**.

IMPROVING EFFICACY

Zinc phosphide bait deteriorates when exposed to rain, humidity and wet acidic soils. The longer the product is exposed to these conditions, the less effective it will be. Efficacy may be improved if applications are made after the soil pH is increased. Avoid treating before morning dew evaporates.

Special Crop Use Restrictions:

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

24(c) Registrant:
HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956