Ramik® green

BAIT PACKS

KILLS RATS, MICE, AND MEADOW VOLES*

Fish Flavored, Weather-Resistant Rodenticide

For Control of Rats, Mice, and Meadow Voles*

Indoors and Outdoors

Active Ingredient:
Diphacinone (2-Diphenylacetyl-1, 3-Indandione).....0.005%
Other Ingredients:.....99.995%
Total.....100.000%

EPA Reg. No. 61282-48 EPA Est. No. 61282-WI-01

This product may only be used inside and within 100 feet of buildings or inside of transport vehicles (ships, trains, or aircraft).

This Product May Not Be Sold in Packaging that Holds Less Than 4 Pounds of Bait.

***Not permitted for use against the following species in California: Cotton rat, Polynesian rat, White-throated woodrat, Southern plains woodrat, Mexican woodrat, Eastern harvest mouse, Golden mouse, Meadow vole**

KEEP OUT OF REACH OF CHILDREN

CAUTION

Net Contents: 16 x 4 oz. (113.4 g)
Net Weight: 4 lbs. (1.81 kg)

NEOGEN®

SPECIMEN LABEL

Ramik green BAIT PACKS

FIRST AID

If Swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor.
If on Skin or Clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor immediately for treatment advice.
If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor immediately for treatment advice.

NOTE TO PHYSICIAN
If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K, intramuscularly or orally repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING
If animal eats bait, call veterinarian or 1-800-498-5743 at once.

Anticoagulant Diphacinone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin time), give Vitamin K.

NOTE TO VETERINARIAN
Keep the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

Personal Protective Equipment (PPE):
All handlers (including applicators) must wear long pants, socks and barrier laminate gloves. Any person who retrieves carcasses or unused bait following application of this product must wear barrier laminate gloves.

User Safety Requirements
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of barrier laminate gloves before removing. As soon as possible, wash hands thoroughly after applying the bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS
This product is extremely toxic to mammals, birds and other wildlife. Dogs, cats and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not allow bait to be exposed on soil surface. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment wash water or rinsate.

Item No. RE491

Manufactured by:
10 Hopkins Drive
Randolph, WI 53956 U.S.A.
NKCCO, Inc. is a wholly-owned subsidiary of Neogen Corporation.

0 23626 00630 0

BA28156-R23

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read the entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

1. Place product in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals or non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be taken from stations when they are filled, units must be secured or otherwise modified. Even tamper proof bait stations are needed in areas open to hooded livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product containers and unused, spoiled, and unconsumed bait as specified on this label.

IMPORTANT: Do not place bait in or near open cribs, pits or other non-target wildlife habitat. Bait stations must be used in non-target areas. Bait stations must be secured or otherwise modified. Even tamper proof bait stations are needed in areas prone to vandalism.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

USE RESTRICTIONS:

This product may only be used to control the following rodent pests in and around man-made structures: House mouse (*Mus musculus*), Norway rat (*Rattus norvegicus*), roof rat (*Rattus rattus*), Cotton rat (*Sigmodon hispidus*)*, Eastern harvest mouse (*Reithrodontomys humulis*)*, Golden mouse (*Ochrotomys nuttalli*)*, Polynesian rat (*Rattus exulans*)*, White-footed woodrat (*Neotoma pennsylvanica*)*, Southern plains woodrat (*Neotoma microtis*)*, Mexican woodrat (*Neotoma mexicana*)*, and Meadow vole (*Microtus pennsylvanicus*)*. This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be inaccessible to commensal rodent reasons and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other premises or temporary residences; food processing facilities; industrial and commercial buildings; farm implements; agricultural and public buildings; transport vehicles (ships, trains, aircraft); docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting beyond 100 feet from a structure as defined above, is prohibited. This product must be applied directly to food or feed crops.

Burrow baiting with Ramik green bait packs is prohibited.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice or meadow voles will most likely find and consume the bait. Generally, these areas are along walls, by grained openings, or at inside corners, or corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Place 1 to 4 bait packs per placement. Space placements at intervals of 15 to 30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 21 days or until there no longer are signs of new feeding by rats.

MICE AND MEADOW VOLES: Open package and apply ¼ to ½ ounce (~2 level tablespoons) of bait at each placement location. Space placements at 6- to 12-foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse or meadow vole activity. Maintain an uninterrupted supply of fresh bait for at least 21 days or until there are no longer signs of new feeding by mice or meadow voles.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using barrier laminate gloves collect and dispose of all dead, much as possible. If infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

***Not permitted for use against the following species in California: Cotton rat, Polynesian rat, White-throated woodrat, Southern plains woodrat, Mexican woodrat, Eastern harvest mouse, Golden mouse, Meadow vole**

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

Pesticide Containers: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration.

Paper Containers: Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration.

DISCLAIMER

DISCLAIMER: To the extent permitted by applicable law, Seller makes no representation or warranty, either express or implied, for results due to misuse, improper handling or improper storage of this material. Nor does Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or improper storage of this material.

For non-emergency (eg., current product information), call: 1-800-621-8829

Tear Notch .625"
Zipper 1.50"
375 Seal Area
14.500"
2.750"
34.500" Web Width
5.500" Bottom Gusset
12.500" Repeat
Seal .375"
Cust: Green Pak
12.5x14.5x5.58G
032104ZNC
Date: 03/04/21