

For Laboratory Use

Store at 2-8°C



Staph Supplement Part #SI-118B

5 mL Vial

(One 5 mL vial contains enough supplement for 50 Staph medium vials.)

Each Vial Contains:

Nalidixic Acid.....	0.0045 gm
Colistin Sulfate.....	0.0045 gm

Usage Instructions:

1. Add 5.0 mL of sterile water to the vial and gently mix. Allow the vial to sit for 10 minutes and gently mix again.
2. Aseptically, add 0.10 mL of the reconstituted supplement to each Soleris Staph test vial containing the sample being tested.
3. The supplement should be used within 7 days following reconstitution, store refrigerated.



620 Leshar Place, Lansing, MI 48912
 800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
 e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2009. Neogen is a registered trademark and Soleris is a trademark of Neogen Corporation.
 C0120504B S-SI118B-0109

For Laboratory Use

Store at 2-8°C



Staph Supplement Part #SI-118B

5 mL Vial

(One 5 mL vial contains enough supplement for 50 Staph medium vials.)

Each Vial Contains:

Nalidixic Acid.....	0.0045 gm
Colistin Sulfate.....	0.0045 gm

Usage Instructions:

1. Add 5.0 mL of sterile water to the vial and gently mix. Allow the vial to sit for 10 minutes and gently mix again.
2. Aseptically, add 0.10 mL of the reconstituted supplement to each Soleris Staph test vial containing the sample being tested.
3. The supplement should be used within 7 days following reconstitution, store refrigerated.



620 Leshar Place, Lansing, MI 48912
 800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
 e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2009. Neogen is a registered trademark and Soleris is a trademark of Neogen Corporation.
 C0120504B S-SI118B-0109

For Laboratory Use

Store at 2-8°C



Staph Supplement Part #SI-118B

5 mL Vial

(One 5 mL vial contains enough supplement for 50 Staph medium vials.)

Each Vial Contains:

Nalidixic Acid.....	0.0045 gm
Colistin Sulfate.....	0.0045 gm

Usage Instructions:

1. Add 5.0 mL of sterile water to the vial and gently mix. Allow the vial to sit for 10 minutes and gently mix again.
2. Aseptically, add 0.10 mL of the reconstituted supplement to each Soleris Staph test vial containing the sample being tested.
3. The supplement should be used within 7 days following reconstitution, store refrigerated.



620 Leshar Place, Lansing, MI 48912
 800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
 e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2009. Neogen is a registered trademark and Soleris is a trademark of Neogen Corporation.
 C0120504B S-SI118B-0109

For Laboratory Use

Store at 2-8°C



Staph Supplement Part #SI-118B

5 mL Vial

(One 5 mL vial contains enough supplement for 50 Staph medium vials.)

Each Vial Contains:

Nalidixic Acid.....	0.0045 gm
Colistin Sulfate.....	0.0045 gm

Usage Instructions:

1. Add 5.0 mL of sterile water to the vial and gently mix. Allow the vial to sit for 10 minutes and gently mix again.
2. Aseptically, add 0.10 mL of the reconstituted supplement to each Soleris Staph test vial containing the sample being tested.
3. The supplement should be used within 7 days following reconstitution, store refrigerated.



620 Leshar Place, Lansing, MI 48912
 800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
 e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2009. Neogen is a registered trademark and Soleris is a trademark of Neogen Corporation.
 C0120504B S-SI118B-0109