

## Listeria Chromogenic Selective Supplement (NCM4002)

### Intended Use

A selective culture media supplement designed for the isolation of *Listeria monocytogenes* according to ISO 11290:2017. Listeria Chromogenic Selective Supplement is intended for use in culture media for the microbiological analysis of food, animal feed, water and environmental samples, and is for use with:

NCM1004      Listeria Chromogenic Agar (Ottaviani & Agosti)

### Typical Formulation

Nalidixic Acid	20.0 mg/L
Ceftazidime	20.0 mg/L
Polymyxin B	76 700 IU/L
Amphotericin	10.0 mg/L

### Typical Formulation by Vial Size

	Per 500 mL Vial NCM4002-0.5
Nalidixic Acid	10 mg
Ceftazidime	10 mg
Polymyxin B	38 350 IU
Amphotericin	5 mg

### Precaution

Refer to SDS

### Preparation

1. Using aseptic technique, remove the crimp cap and rubber stopper. Aseptically retain the stopper.
2. Reconstitute the contents of the vial using the appropriate volume of sterile deionized/RO water as indicated in the table below.
3. Replace the rubber stopper and gently shake or invert the contents to dissolve.
4. Add the vial contents to the cooled medium as per the label directions.

	NCM4002-0.5
Supplement Reconstitution Volume	5 mL
Media Prep Volume	500 mL

### Quality Control Specifications

Lyophilized Appearance: Yellow/orange pellet which may break up with agitation.

Reconstituted Appearance: Cloudy, yellow solution.

### Storage

Store sealed vial at 2-8°C away from direct sunlight.

### Expiration

Refer to expiration date stamped on the container. Expiry applies to unopened dehydrated supplement.



# Technical Specification Sheet



## **Precautions**

Discard if lyophilized supplement demonstrates any moisture uptake or appears to have liquefied; discard if reconstituted supplement demonstrates any precipitate.

Revision: 0 Effective Date: 10/7/2020



620 Lesher Place • Lansing, MI 48912  
800-234-5333 (USA/Canada) • 517-372-9200  
foodsafety@neogen.com • foodsafety.neogen.com