

Preston Supplement (NCM4038)

Intended Use

A selective culture media supplement for the isolation of *Campylobacter* spp. Preston Supplement is intended for use in culture media for the microbiological analysis of food, animal feed, water and environmental samples, and is for use with:

NCM0189 Nutrient Broth No. 2

Typical Formulation

Rifampicin	10.0 mg/L
Polymyxin B	5000 IU/L
Trimethoprim	10.0 mg/L
Amphotericin B	10.0 mg/L

Typical Formulation by Vial Size

	Per 500 mL Vial NCM4038-0.5
Rifampicin	5.0 mg
Polymyxin B	2500 IU
Trimethoprim	5.0 mg
Amphotericin B	5.0 mg

Precaution

Refer to SDS

Preparation

1. Using aseptic technique, remove the crimp cap and rubber stopper. Aseptically retain the stopper.
2. Reconstitute the contents of the vial using the appropriate volume of sterile 50% ethanol as indicated in the table below.
3. Replace the rubber stopper and gently shake or invert the contents to dissolve.
4. Add the vial contents to the cooled medium as per the label directions.

	NCM4038-0.5
Supplement Reconstitution Volume	5 mL
Media Prep Volume	500 mL

Quality Control Specifications

Lyophilized Appearance: Homogenous red dried pellet; may break up with agitation.

Reconstituted Appearance: Red opaque solution without a precipitate.

Storage

Store sealed vial at 2-8°C away from direct sunlight.

Expiration

Refer to expiration date stamped on the container. Expiry applies to unopened dehydrated supplement.

Technical Specification Sheet



Precautions

Discard if lyophilized supplement demonstrates any moisture uptake or appears to have liquefied; discard if reconstituted supplement demonstrates any precipitate.

Revision: 0 Effective Date: 11/16/2020



620 Lesher Place • Lansing, MI 48912
800-234-5333 (USA/Canada) • 517-372-9200
foodsafety@neogen.com • foodsafety.neogen.com