

AquaPrime®

PERASIDE™ 15

AquaPrime® Peraside™ 15 is a peroxyacetic acid-based microbiocide developed for Equipment Sanitizing, Disinfection, Aseptic Packaging and Bacteria, Fungi, Slime and Odor Control in: Fruit and Vegetable Process Water Systems

Active Ingredient:	
Peroxyacetic Acid.....	15.0%
Hydrogen Peroxide.....	22.0%
Inert Ingredients:	63.0%
Total:	100.0%

EPA Registration No: 63838-2-66171

EPA Est. No. 63838-CA-01: 63838-AR-001

Before Using This Product, Please Read This Entire Label Carefully.

**KEEP OUT OF REACH OF CHILDREN
DANGER-PELIGRO**



Certified to NSF/ANSI 60

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If in Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

QUESTIONS? 1-209-581-9576

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER CORROSIVE. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles, face shield, rubber gloves and protective clothing with long sleeves when handling. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse. Do not enter an enclosed area without proper respiratory protection, or when uncoupling of product transfer hoses. Wear a minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and combination N1, R, or P filters; or a NIOSH-approved gas mask with OV canisters; or a NIOSH-approved powered air purifying respirator with OV cartridges and combination HE filters when handling concentrate product.

PHYSICAL OR CHEMICAL HAZARDS:

STRONG OXIDIZING AGENT. CORROSIVE: Mix only with water below 140° F. Product must be diluted in accordance with label directions prior to use. This product is not combustible; however, at temperatures exceeding 156° F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to birds, fish and aquatic invertebrates. Caution must be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of the National Pollution Discharge System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage plant authority.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note: All volumes given in ounces are fluid ounces.

SANITIZATION

This peroxyacetic acid sanitizer is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, filters, evaporators, pasteurizers, and aseptic equipment in dairies, breweries, wineries, beverage and food processing/packing plants, and egg processing/packing equipment surfaces. This product is effective as a sanitizer when solution is prepared in water of up to 400 ppm hardness as CaCO₃. This product has demonstrated greater than 99.999% reduction of Staphylococcus aureus and Escherichia coli in the AOC Germicidal and Detergent Sanitizing Action of Disinfectants study.

Sanitizing Food Contact Surfaces: Sanitize with a concentration of 0.7-3.8 fl. oz. of this product diluted in 10 gallons of water (93-500 ppm active peroxyacetic acid and 136-733 ppm active hydrogen peroxide). Use immersion, spray or circulation techniques as appropriate to the equipment. All surfaces must be exposed to sanitizing solution for a period of at least 60 seconds or more if specified by a governing code. Drain thoroughly and allow to air dry. Do not rinse.

Sanitization of Conveyors and Equipment for Meat, Poultry, Seafood, Dairy, Fruit, Nuts and Vegetables: This product is effective against the gram positive organism Staphylococcus aureus and gram negative organism Escherichia coli. For use in the static or continuous sanitizing, washing or rinsing of conveyors, slicers, saws, and equipment, apply a solution of this product using a recommended 0.7-3.8 fl. oz. per 10 gallons of water (93-500 ppm active peroxyacetic acid and 136-733 ppm active hydrogen peroxide). Apply sanitizer solution to the return portion of the conveyor or equipment using spray or similar means of wetting surfaces, so as to prevent puddling. Allow sanitizer to thoroughly wet surface for a minimum 60 seconds contact time. No rinse is needed.

Final Bottle or Container Rinse: This product may be used as a final sanitizer rinse for pre-cleaned returnable and non-returnable bottles or containers at 93-500 ppm active peroxyacetic acid and 136-733 ppm active hydrogen peroxide (0.7-3.8 fl. oz. of this product diluted in 10 gallons of water). The container must be drained as much as is practical prior to filling operations.

POULTRY, SWINE, LIVESTOCK WATERING OPERATING SYSTEMS

After watering lines have been cleaned, use this product at 0.3-42 fl. oz. per 100 gallons of water (4-559 ppm as peroxyacetic acid) to control algae and bacteria in drinking water and to control mineral build up in watering lines. Stop the use of this product twenty-four (24) hours prior to vaccination via the water line.

ANIMAL PREMISES

This product is designed for use in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries, and livestock quarters. When used as directed, this product is specifically designed to disinfect, deodorize and clean inanimate, hard, surfaces such as walls, floors, sink tops, furniture, operating tables, kennel runs, cages and feeding equipment. In addition, this product will deodorize those areas which are generally difficult to keep smelling fresh, such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

Disinfection of Poultry Premises, Trucks, Coops and Crates: For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use. Remove all poultry and feeds from premises, trucks, coops and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.16-0.61% v/v (1.0-3.9 fl. oz. per 5gal) solution of this product for a period of 10 minutes. This is equivalent to 266-1036 ppm PAA and 390-1520 ppm H₂O₂. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waters with a detergent and rinse with potable water before reuse. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse. See your technical representative for specific recommendations for all cleaning and rinsing requirements.

Disinfection and Deodorizing of Animal Housing Facilities (Barns, Kennels, Hutches, Etc.): Remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering equipment. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces by applying a 0.16-0.61% v/v (1.0-3.9 fl. oz. per 5 gal) solution of this product with a mop, brush or spray. This is equivalent to 266-1036 ppm PAA and 390-1520 ppm H₂O₂. Wet all surfaces and allow to remain wet for 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. Ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

Commercial Agricultural or Industrial Water System Injection for the Oxidation of Odors When They Form: In well systems, inject this product with a peristaltic pump through a ¼ inch plastic tube (or other comparable injection system) inserted into the well all the way to the well intake screens. Test the most distant faucet a hydrogen peroxide test strip until a residual of 20-25 ppm hydrogen peroxide is detected indicating that the system has been thoroughly treated. Adjust the injection pump to deliver 25 ppm on a consistent basis to the water system. When treating the watering system for sick animals, test the most distant faucet with a hydrogen peroxide test strip until a residual of 100-1,000 ppm hydrogen peroxide is detected indicating that the system has been thoroughly treated. When this level is reached immediately reduce the injection until the test strip indicates the 25 ppm hydrogen peroxide is reached and the system is maintained at the 25 ppm level. This process can be repeated monthly to ensure a clean water system and avoid screen clogging.

Cleaning Poultry and Livestock Drinking Water Lines:

For drinking water lines using holding tanks make a stock solution by one of the following methods:

- Drinking water lines 500 feet in length or less: mix 0.78 gallons (100 fl. oz.) of this product with 100 gallons of water.
- Drinking water lines exceeding 500 feet in length: mix 1.55 gallons (200 fl. oz.) of this product with 200 gallons of water.

Pump the stock solution, completely filling the drinking water lines.

If the drinking water lines are not supplied by water from holding tanks, prepare a stock solution by one of the following methods:

- Mix 0.38 gallons (49 fl. oz.) of this product with 49 gallons of water in a 50 gallon tank, pumping this solution into the water line, repeating the process as often as needed, until water line is filled.
- Fill the water line, using a proportioner, set to inject this product undiluted at a rate of 1:11 (0.9%).

After the waterline is filled with the stock solution, activate nipple drinkers to ensure contact with drinkers.

Allow the stock solution to remain in the water lines for 24-48 hours.

Flush lines with fresh water until water is visibly clear.

Always make a fresh stock solution before use.

TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS

This product can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables for the control of spoilage and decay causing bacteria and fungi in commercial operations and packinghouses.

Batch, Continuous or Spray System Processes: Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method. Add this product to no more than 500 ppm residual peroxyacetic acid to the use solution in accordance with Food Contact Notification #1738, effective March 28, 2017. This can be accomplished by initially adding 3.8 fl. oz. per 10 gallons of water. The recommended concentration is between 30-300 ppm as peroxyacetic acid (0.23-2.3 fl. oz. per 10 gallons of water). The final concentration necessary to accomplish the intended task will vary from plant-to-plant. The fruits and vegetables can be continuously sprayed or submerged (dipped) in the resulting solution. Periodic or continuous additions of this product to maintain the required concentration may be added as necessary. It is also recommended to apply this product during the washing, chilling, or physical cleaning processes, including the roller-spreader, washer or brush washer manifold, dip tank, or sorting processes. Contact time of 60 seconds is recommended to insure efficacy. A potable water rinse is not required.

Fogging: (Not for Use in California): For raw agricultural commodities, commercially-applied fogging methods may be used provided the dilution rates of the resultant solution does not exceed those prescribed in this section (3.8 fl. oz. per 10 gal. of water). A potable water rinse is not required. Conventional corrosion-resistant fogging devices are recommended. Vacate the area of all personnel prior to, during and after fogging until the total peroxide concentration is below 1.0 ppm, or there is no strong odor present, characteristic of acetic acid.

STORAGE AND DISPOSAL

Storage: Never return this product to the original container after it has been removed. Avoid all contaminants, especially dirt, caustic, reducing agents, and metals. Contamination and impurities will reduce shelf life and can induce decomposition. In case of a decomposition, isolate container, spray container with cool water and dilute this product with large volumes of water. Avoid damage to containers. Keep container closed at all times when not in use. **Keep container out of direct sunlight.** To maintain product quality, store at temperatures below 86°F.

Procedure for Leak or Spill: Stop leak if this can be done without risk. Shut off ignition sources: no flames, smoking, fires, or spark producing tools. Keep combustible and organic materials away. Flush spilled material with large quantities of water. Undiluted material must not enter confined spaces.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance. If material has been spilled, an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all local, state and Federal environmental laws, rules, regulations, standards, and other requirements. Because acceptable methods of disposal may vary by location, regulatory agencies must be contacted prior to disposal. This product which is to be discarded, must be disposed of as hazardous waste after contacting the appropriate local State or Federal agency to determine proper procedures.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Offer for recycling, if available. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Container Handling: (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat the procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

24 hr Emergency ChemTel Number: 1-800-424-9300

DOT: UN 3109, Organic Peroxide Type F, Liquid (<=25% Peracetic Acid with <=26% Hydrogen Peroxide) 5.2 (8)

Lot # _____



Manufactured by: Preserve International®
944 Nandino Blvd., Lexington, KY 40511 U.S.A.
L6541-1020

Preserve International® is a wholly-owned subsidiary of Neogen® Corporation

Item No. 480020/2.5



7 89703 00492 7



Net Contents: 2.5 gal (9.46 L)