

# Judgement

## *soft bait*



**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

**KILLS RATS, MICE AND MEADOW VOLES**

### Agricultural Building Use

For Use In and Around Agricultural Buildings Only

**FOR INDOOR AND OUTDOOR USE**

**KILLS RATS, MICE AND MEADOW VOLES**

**KILLS WARFARIN RESISTANT NORWAY RATS**

See Back Panel for additional precautionary labeling

#### Active Ingredient:

Difenacoum (CAS #56073-07-5).....0.005%

Other Ingredients.....99.995%

Total.....100.000%


^Contains Denatonium Benzoate

EPA Reg. No. 87235-4-61282

EPA Est. No. 87235-GBR-001

NOT REGISTERED FOR SALE  
OR USE IN CALIFORNIA

L6979-0520



#### FIRST AID

**If swallowed:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**If on skin or clothing:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor immediately for treatment advice.

**If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

#### Note To Physician

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

#### Treatment For Pet Poisoning

If animal eats bait, call veterinarian at once.

#### Note To Veterinarian

Anticoagulant Difenacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.



**Net Weight:**  
22 lbs. (9.98 kg)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers (including applicators) must wear: shoes plus socks, long pants and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**USER SAFETY REQUIREMENTS:** Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

### RESTRICTIONS:

1. Do not expose children, pets, or non-target animals to rodenticides.
2. Store unused product out of reach of children, pets and non-target animals.
3. Bait stations are mandatory for outdoor, above ground use. Tamper resistant bait stations must be used if children, pets, domestic animals, non-target animals or birds may access the bait.
4. Apply bait indoors in locations out of reach of children, pets, domestic animals and non-target animals, or in tamper-resistant bait stations. These bait stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
5. Do not apply bait directly to food of feed crops
6. Do not broadcast bait.
7. Do not apply to burrows
8. Do not sell this product in individual containers holding less than 8 pounds of bait.
9. Fence and perimeter baiting, beyond 100 feet from a structure is prohibited.
10. Not registered for sale or use in California.

This product can only be used to control Norway rats (*Rattus norvegicus*), Roof rats (*Rattus rattus*), House mice (*Mus musculus*), Cotton rat (*Sigmodon hispidus*), Eastern harvest mouse (*Reithrodontomys humulus*), Golden mouse (*Ochrotomys nuttalli*), Meadow vole (*Microtus pennsylvanicus*), Mexican woodrat (*Neoto-*

*ma Mexicana*), Polynesian rat (*Rattus exulans*), Southern plains woodrat (*Neotoma micropus*) and White throated woodrat (*Neotoma albigula*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice or meadow voles will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats, mice or meadow voles have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each bag of Judgement soft bait contains approximately 0.54 ounces (15.00 grams) of bait.

**RATS:** Apply 7 to 26 bait bags per placement for controlling rats. Space placements 15 to 30 feet apart at appropriate locations within the infested area. Maintain a continuous supply of fresh bait for at least 10 days or until signs of new feeding by rats on bait cease to appear.

**MICE & MEADOW VOLE:** Apply 2 bait bags per placement usually spaced 8 to 12 feet apart, reducing to every 4 to 6 feet at points of very high mouse or meadow vole activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse or meadow vole activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

**DISCLAIMER:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

For non-emergency (e.g., current product information), call: 1-800-621-8829

**Manufactured for:** HACCO, Inc.

110 Hopkins Drive  
Randolph, WI 53956 U.S.A.

HACCO, Inc. is a wholly-owned subsidiary  
of Neogen® Corporation  
L6979-0520

EPA Reg. No. 87235-4-61282

EPA Est. No. 87235-GBR-001

Item No. 373424

